

Biorbyt Ltd. 5 Orwell Furlong, Cowley Road, Cambridge CB4 0WY, United Kingdom Email: info@biorbyt.com Phone: +44 (0)1223 859 353 | Fax: +44 (0)1223 280 240 Biorbyt LLC. 369 Pine Street, Suite #103, San Francisco California, 94104, United States Email: info@biorbyt.com Phone: +1 (415) 906 5211 | Fax: +1 (415) 651 8558

Product Datasheet

TLR4 antibody orb152207

Description	Rabbit polyclonal to TLR4 (RPE). The Toll-like receptor (TLR) family in mammal comprises a family of trans-membrane proteins characterized by multiple copies of leucine rich repeats in the extracellular domain and 1L-1 receptor motif in the cytoplasmic domain. Like its counterparts in Drosophila, TLRs signal through adaptor molecules. The TLR family is a phylo-genetically conserved mediator of innate immunity that is essential for microbial recognition. Ten human homologs of TLRs (TLR1-10) have been described. Amount this family of receptors, TLR2 and TLR4 have been most studied. These studies have suggested that TLR2 and TLR4 may serve as potential main mediators of LPS signaling (4,5). The mouse TLR4 cDNA codes for a protein consisting of 839 amino acids, with an approximate molecular weight of 90kDa
Species/Host	Rabbit
Reactivity	Human, Mouse
Conjugation	RPE
Tested Applications	FACS, FC, ICC, IHC, WB
Immunogen	Developed against a synthetic peptide corresponding to amino acids 420-435 of human TLR4
Target	TLR4
Concentration	1 mg/ml
Preservatives	PBS, 50% glycerol, 0.09% sodium azide *Storage buffer may change when conjugated
Storage	Conjugated antibodies should be stored at 4°C
Note	For research use only.
Clonality	Polyclonal
MW	75-80kDa
Uniprot ID	O00206
NCBI	7099, NP_612564.1
Entrez	7099
Dilution Range	WB (1:500), IHC (1:50)
Expiration Date	12 months from date of receipt.
Application Notes	2 μ g/ml of SPC-200 was sufficient for detection of TLR4 in 100 ng of partial recombinant mouse TLR4 protein by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.