

Biorbyt Ltd.

5 Orwell Furlong, Cowley Road, Cambridge CB4 0WY, United Kingdom Email: info@biorbyt.com

Phone: +44 (0)1223 859 353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

369 Pine Street, Suite #103, San Francisco California, 94104, United States Email: info@biorbyt.com

Phone: +1 (415) 906 5211 | Fax: +1 (415) 651 8558

Product Datasheet

Covid-19 & SARS-CoV Nucleoprotein antibody orb669768

Description Mouse monoclonal antibody to Covid-19 & SARS-CoV Nucleoprotein

Species/Host Human

Reactivity Virus

Conjugation Unconjugated

Tested Applications ELISA, IF

Immunogen The original antibody was generated by cloning the variable regions of the

scFvs selected from phage display libraries into separate vectors for IgG1 heavy-chain and light-chain expression. The harvested supernatents were then purified on protein A columns. The original immunogen was the whole

irradiated virion.

Target Covid-19 & SARS-CoV Nucleoprotein

Concentration 1 mg/ml

Preservatives PBS with 0.02% Proclin 300.

Storage Store at 4°C for up to 3 months. For longer storage, aliquot and store at

-20°C.

Note For research use only.

Isotype IgG1

Clonality Monoclonal

Purity Purified

Clone ID CR3018 (03-018)

Uniprot ID P59595, P0DTC9

Expiration Date 12 months from date of receipt.

Application Notes This antibody is recommended for detection of SARS CoV2 protein N (nucleoprotein). This

antibody binds both the nucleocapsid protein of the SARS-CoV and SARS CoV-2 (2019-nCoV). Initial characterization of the antibody for binding to 2019-nCoV was done using ELISA. This antibody shows potential to be used for development of diagnostic assays. Various isotype versions of the antibody namely human IgG1, IgG3, IgM, IgA and the less common IgG2 and IgG4 are available for the investigation of their role in response to SARS CoV2. Competitive ELISA of this antibody with CR3009 suggests that both these antibodies bind different epitopes of the N protein of SARS CoV. Thus, a combination of these two antibodies is suggested for virus capture assays. Immunofluorescence staining was used to demonstrate binding of CR3018 to SARS-CoV infected Vero cells.

(PMID:15650189)